

STATE-OF-THE-ART PAPER

# Heart Failure Clinical Trials in East and Southeast Asia



## Understanding the Importance and Defining the Next Steps

Robert J. Mentz, MD,<sup>a</sup> Lothar Roessig, MD,<sup>b</sup> Barry H. Greenberg, MD,<sup>c</sup> Naoki Sato, MD, PhD,<sup>d</sup> Kaori Shinagawa, MD, PhD,<sup>e</sup> Daniel Yeo, MBBS,<sup>f</sup> Bernard W.K. Kwok, MBBS,<sup>g</sup> Eugenio B. Reyes, MD,<sup>h</sup> Henry Krum, MBBS, PhD,<sup>i,†</sup> Burkert Pieske, MD,<sup>j</sup> Stephen J. Greene, MD,<sup>a</sup> Andrew P. Ambrosy, MD,<sup>a</sup> Jacob P. Kelly, MD,<sup>a</sup> Faiez Zannad, MD,<sup>k,l,m,n,o</sup> Bertram Pitt, MD,<sup>p</sup> Carolyn S.P. Lam, MBBS<sup>q</sup>

### ABSTRACT

Heart failure (HF) is a major and increasing global public health problem. In Asia, aging populations and recent increases in cardiovascular risk factors have contributed to a particularly high burden of HF, with outcomes that are poorer than those in the rest of the world. Representation of Asians in landmark HF trials has been variable. In addition, HF patients from Asia demonstrate clinical differences from patients in other geographic regions. Thus, the generalizability of some clinical trial results to the Asian population remains uncertain. In this article, we review differences in HF phenotype, HF management, and outcomes in patients from East and Southeast Asia. We describe lessons learned in Asia from recent HF registries and clinical trial databases and outline strategies to improve the potential for success in future trials. This review is based on discussions among scientists, clinical trialists, industry representatives, and regulatory representatives at the CardioVascular Clinical Trialist Asia Forum in Singapore on July 4, 2014. (J Am Coll Cardiol HF 2016;4:419-27)  
© 2016 by the American College of Cardiology Foundation.

**H**eat failure (HF) is a major public health problem worldwide (1,2). In Asia, aging populations and large increases in cardiovascular risk factors have contributed to a high burden of HF (3). HF patients from Asia differ in clinical characteristics from patients elsewhere and yet have similarly poor or even worse outcomes than

HF patients from the West (Table 1) (4). Given the diversity of countries in Asia (Online Appendix), clinical phenotypes and practice patterns vary widely, just as practice patterns vary across Europe and the Americas.

The last several decades have seen therapeutic advances for HF patients with reduced ejection

From the <sup>a</sup>Duke Clinical Research Institute and the Department of Medicine, Division of Cardiology, Duke University Medical Center, Durham, North Carolina; <sup>b</sup>Bayer HealthCare, Wuppertal, Germany; <sup>c</sup>Division of Cardiology, University of California San Diego, San Diego, California; <sup>d</sup>Internal Medicine, Cardiology, and Intensive Care Medicine, Nippon Medical School Musashi-Kosugi Hospital, Kawasaki, Kanagawa, Japan; <sup>e</sup>Office of New Drug II, Pharmaceuticals and Medical Devices Agency, Chiyoda-ku, Tokyo, Japan; <sup>f</sup>Gleneagles Hospital, Singapore; <sup>g</sup>Farrer Park Medical Centre, Singapore; <sup>h</sup>Cardiology, Manila Doctors Hospital, Manila, Philippines; <sup>i</sup>Centre of Cardiovascular Research and Education in Therapeutics, School of Public Health and Preventive Medicine, Monash University, Melbourne, Australia; <sup>j</sup>Department of Cardiology, Charité University Medicine, Campus Virchow Klinikum, and German Heart Center Berlin, Berlin, Germany; <sup>k</sup>INSERM, Centre d'Investigations Cliniques-Plurithématique 1433, Vandoeuvre-lès-Nancy, France; <sup>l</sup>INSERM U1116, Nancy, France; <sup>m</sup>Université de Lorraine, Nancy, France; <sup>n</sup>CHU Nancy, Pôle de Cardiologie, Institut Lorrain du Cœur et des Vaisseaux, Vandoeuvre-lès-Nancy, France; <sup>o</sup>F-CRIN INI-CRCT (Cardiovascular and Renal Clinical Trialists), Nancy, France; <sup>p</sup>University of Michigan School of Medicine, Ann Arbor, Michigan; and the <sup>q</sup>National Heart Centre Singapore and Duke-National University of Singapore, Singapore. Dr. Mentz is supported by National Institutes of Health grant U10HL110312. Dr. Roessig is an employee of Bayer HealthCare. Dr. Greenberg is a consultant for Novartis. All other authors have reported that they have no relationships relevant to the contents of this paper to disclose.

<sup>†</sup>Dr. Henry Krum died November 28, 2015.

Manuscript received December 3, 2015; revised manuscript received December 22, 2015, accepted January 8, 2016.

## ABBREVIATIONS AND ACRONYMS

**ACEI/ARB** = angiotensin-converting enzyme inhibitor/angiotensin receptor blocker

**CAD** = coronary artery disease

**GDMT** = guideline-directed medical therapy

**HFREF** = heart failure with reduced ejection fraction

**ICD/CRT** = implantable cardioverter-defibrillator/cardiac resynchronization therapy

**LOS** = length of stay

**MRA** = mineralocorticoid receptor antagonist

fraction (EF) (5), including recent trials of sacubitril/valsartan (6) and ivabradine (7) therapy. Older trials either did not include Asians or included small numbers of patients of Asian ethnicity from Western countries. Given the need to enroll large numbers to demonstrate outcome benefits as well as recent challenges with enrollment and cost in North American and certain European countries, contemporary trials have enrolled globally (8). Representation of patients from Asia was relatively low in many previous trials due to perceived and/or actual challenges to generalizability and trial infrastructure and conduct. Regulatory approval of HF drugs in Asian countries has largely relied on data extrapolated from Western

populations. More recently, Asian regulatory authorities have been requiring that study populations include Asians from Asia (9) in order to support approval.

We reviewed differences between HF patients from East and Southeast Asia (Online Appendix) and compared them to HF patients in the rest of the world. We focus on East and Southeast Asia because these regions within Asia have more robust registry and trial data available to date. We summarized observations within the context that heterogeneity exists even among regions within Asia. We describe lessons learned from HF datasets and outline strategies to improve future trials. This review is based on discussions among scientists, trialists, industry representatives, and regulators at the Cardiovascular Clinical Trialist Asia Forum in Singapore on July 4, 2014. To identify additional relevant published data not discussed, we searched MEDLINE for articles published between January 1994 and December 2015 (see the Online Appendix for the search strategy).

## BURDEN OF HF

Limited data are available regarding the true incidence and prevalence of HF in Asia (3). Studies of hospitalized patients in Singapore, Malaysia, and Taiwan found that 3% to 7% of admissions were due to HF in the 1990s to early 2000s (10-12). In a community-based survey, the prevalence of HF in China among populations 55 to 74 years of age was 1.3%, with an estimated overall adult HF population >4 million (13). Similarly, in Japan, it is estimated that 1 million people have HF (14), which equates to a prevalence of ~1%. Although prevalence estimates in the general population are lower in Asia than in the West (1,15), this translates to a higher absolute burden

of disease in Asia because of larger population sizes. For example, even with conservative estimates of HF prevalence, the absolute number of individuals with HF in Asia is >20 million (16). Regarding HF hospitalizations, in Singapore there was a 38% increase from 1991 to 1998, which is approximately 5% per year (10). This matches the 5% yearly increase in all-cause hospitalizations from 2004 to 2012. However, in recent years, HF hospitalizations have been rising at 10% each year (17). Furthermore, the at-risk population is increasing at a faster rate in Asia than in other parts of the world, with aging of the population and increases in the prevalence of coronary artery disease (CAD), tobacco use, diabetes, and obesity. For instance, in 2007, there were 305,700 people older than 65 years of age in Singapore (6.7% of the population). This increased more than 30% to 404,500 in 2013 (7.5% of the population) (18). Thus, the burden of HF in Asia is expected to increase and be comparatively larger than that in the West over subsequent decades (14).

## HF PHENOTYPE


Data from trials and registries in Asia provide insights into the profile of HF in this region (Central Illustration). ADHERE-AP (Acute Decompensated Heart Failure Registry-Asia Pacific) was an acute HF registry that included 10,171 patients hospitalized with HF from 8 Asia-Pacific countries (Singapore, Thailand, Indonesia, Australia, Malaysia, the Philippines, Taiwan, and Hong Kong). HF patients in Asia-Pacific were younger than those from other regions. The median age was 67 to 70 years in Asia versus 70 to 75 years in the United States and Europe (4,19). Moreover, there was variation within different Asia-Pacific countries with the median age at presentation ranging from 53 years in the Philippines to 77 years in Australia and Hong Kong. These differences may be due, in part, to variations in risk factor profile, comorbidity burden, life expectancy, and standards of living (4). Thus, there may be nearly as much heterogeneity and regional variation within Asia-Pacific as between this geographic region and other world regions.

**HF CAUSES.** Compared with other regions where >50% of the population has ischemic HF, there is a lower prevalence of ischemic cardiomyopathy in Asia. For instance, in the ATTEND registry (Acute decompensated heart failure syndromes) consisting of 4,841 acute HF patients enrolled in Japan, 31% of patients had ischemic, 19% had valvular, 18% had hypertensive, and 32% with “other/unspecified” causes (20). A chronic HF registry of 1,078 Japanese patients

**TABLE 1 Comparison of Clinical Characteristics and Outcomes in Asia to Other Regions**

	CHART-1 (19)	CHART-2 (19)	JCARE-CARD (69)	ATTEND (20)	KorAHF (31)	Tseng (12)	ADHERE-AP (4)	ADHERE (30)	OPTIMIZE-HF (70,71)	EHFS II (72)
	2000-2004	2006-2010	2004-2005	2007-2011	2011-2012	2005	2006-2008	2001-2004	2003-2004	2004-2005
	Japan	Japan	Japan	Japan	Korea	Taiwan	AP Region	United States	United States	Europe
	Stage C/D	Stage C/D	AHF	AHF	AHF	AHF	AHF	AHF	AHF	AHF
	N = 1,078	N = 4,735	N = 2,549	N = 4,841	N = 2,066	N = 2,692	N = 10,171	N = 105,388	N = 48,612	N = 3,580
	26 hospitals	24 hospitals	164 hospitals	52 hospitals	10 centers	insured patients	43 hospitals	274 hospitals	259 hospitals	133 hospitals
Age, yrs	69 ± 13	69 ± 12	71 ± 13	73 ± 14	69 ± 14	73 ± 13	67	72 ± 14	73 ± 14	70 ± 13
Males	65%	68%	60%	58%	55%	55%	57%	48%	48%	61%
Ischemia cause	26%	47%	32%	31%	38%	32%	50%	57%	46%	54%
LVEF ≥50%	51/51%	57/69%	42 ± 18	-	40 ± 18	-	53% (LVEF <40%)	34 ± 16/46% (>40%)	39 ± 18	38 ± 15/34% (LVEF ≥45%)
Diabetes	20%	23%	30%	34%	36%	28%	45%	44%	25%	33%
Atrial fibrillation	42%	31%	35%	40%	27%	-	24%	31%	31%	39%
Renal dysfunction	50%	47%	12%	-	-	13%	22%	30%	20%	17%
BMI, kg/m <sup>2</sup>	23.0 ± 3.7	23.8 ± 3.9	22.4 ± 4.1	-	-	-	-	-	-	26.8
SBP, mm Hg	126 ± 19	126 ± 19	117 ± 18	146 ± 37	136 ± 31	-	57% with SBP 90-140	144 ± 33	143 ± 33	135 (110-160)
HR, beats/min	75 ± 14	72 ± 15	70 ± 12	99 ± 29	91 ± 26	-	-	-	87 ± 22	95 (77-114)
Creatinine/eGFR	-/61 ± 31	-/61 ± 24 (stage C)	1.4/52 ± 25	1.4 ± 1.6	1.5 ± 1.6	-	>1.5 mg/dl in 41%	1.8 ± 1.6	1.8 ± 1.8	-
BNP/NT-proBNP, pg/mL	273 ± 353	191 (C), 454 (D)	375 ± 474	707 (362-1,284)	-	-	-	840 (430-1,730)	800 (403-1,660)	-
Chronic HF therapies										
ACEI/ARB	57%/13%	45%/32%	37%/44%	31%/46%	65%	51%	63%	41%/12%	40%/12%	80%
Beta-blocker	28%	49%	49%	67%	44%	25%	41%	48%	53%	61%
MRA	~21%	~22%	42%	-	40%	-	31%	20%	7%	48%
Digoxin	48%	24%	31%	-	24%	32%	34%	28%	23%	31%
CRT/ICD	1.5%	2.9% (C), 15.8% (D)	1.6%/2.0%	2.3%/3.4%	1.3%/1.4%	-	-/1.6%	-	5%	9.1% pacemaker
Inpatient therapies										
IV diuresis	-	-	-	76%	72%	76% (all diuretic)	85%	92%	-	84%
IV vasodilation	-	-	-	78%	40%	-	14% (IV nitrate)	9%	14%	38%
IV inotrope therapy	-	-	-	19%	32%	-	15%	15%	7%	>11%
In-hospital mortality	-	-	3.9% (rEF) 6.5% (pEF)	6.4%	5.2%	3.9%	4.8%	4.0%	3.8%	6.7%
Length of stay, days	-	-	36 (rEF) 31 (pEF)	30 ± 39/21 (14-32 days)	8	15.8 ± 42.7	6.0	4.3	6.4	9 (6-14 days)
30-day mortality	-	-	-	-	1.2%	-	-	-	8.6% at 60-90 days	-
Short-term rehospitalization	-	-	-	-	6% HF rehospitalization at 30 days	-	-	-	29.6% at 60-90 days	-

Values are mean ± SD, median (IQR), or %. If details were not provided in the primary manuscript, this is represented by a "-" sign. If multiple relevant details are provided in the same section, these are separated with a "/" symbol.  
 ACEI = angiotensin-converting enzyme inhibitor; AHF = acute heart failure; ARB = angiotensin receptor blocker; BMI = body mass index; BNP = brain natriuretic peptide; CRT = cardiac resynchronization therapy; GFR = glomerular filtration rate; HR = heart rate; ICD = implantable cardioverter-defibrillator; IV = intravenous; LVEF = left ventricular ejection fraction; MRA = mineralocorticoid receptor antagonist; pEF = preserved ejection fraction; rEF = reduced ejection fraction; SBP = systolic blood pressure.



reported that ischemia was the underlying cause in 26% of cases (19). These observations are notable given the high prevalence of CAD risk factors, including >40% of the Asian population with a smoking history and 45% with diabetes. Younger age may partially explain the discordance between risk factor burden and ischemic prevalence. Recent data suggest that ischemia-driven HF is increasing in Asia. For instance, in Japan, the prevalence of CAD increased from 26% to 47% from 2000 to 2010 (19).

**COMORBIDITIES.** Atrial fibrillation (AF) and diabetes were previously less common in Asia than in other regions, but current trends suggest that both are increasing. Historically, the incidence of AF was lower in the Asian population (21) than in individuals in North America and Western Europe (22). However, from 2001 to 2012, there was a >20-fold increase in AF incidence in China (21). These observations are likely due to the aging of the population as well as comorbidities such as rheumatic heart

disease, lung disease, and diabetes (21). The recent increase in AF incidence in Asia is markedly higher than in North America/Europe (23,24), suggesting that AF may play an even more critical role in HF in Asia in the future (25). Importantly, these observations may also be related to increased disease ascertainment in Asia.

Similarly, diabetes in Asia has increased in recent years due to changes in lifestyle involving physical inactivity and diet changes (26). For instance, in Malaysia, obesity increased from 12% to 15% and diabetes increased from 12% to 15% from 2003 to 2011 (27). Regional changes in lifestyle and diet were strongly associated with diabetes. Similar findings have been seen in other Asian countries (28). Data from the International Diabetes Federation indicate that the Western Pacific and South East Asia regions have the highest levels of diabetes at 138 and 72 million, respectively (29); these numbers are expected to increase more than 40% by 2035. China and India have the largest current and projected populations of people living with diabetes, at 98 and 65 million, respectively, in 2013, growing to 143 and 109 million, respectively, by 2035. Thus, diabetes and obesity will likely become increasingly prevalent in the Asian HF population, which has important implications on clinical management. Other cardiovascular and noncardiovascular comorbidities such as sleep disordered breathing, renal dysfunction, lung disease, depression, and frailty also influence HF management in Asia but are less well characterized than in other world regions.

## HF MANAGEMENT

In-hospital management of acute HF in ADHERE-AP included intravenous (IV) diuretic therapy in 85% of patients and IV inotrope therapy in 15%. IV diuretic use was comparable to North America and Western Europe (~85% to 90%); however, IV inotrope use tends to be higher in Asia than in the United States (4). Similarly, the Japanese ATTEND registry showed high IV inotrope use (19%) and frequent use of IV vasodilators (>70%) (20) compared with only 10% to 15% vasodilator use in the United States (30). IV vasodilator use was also higher in Korea (40%) (31). These variations in practice patterns were observed despite a relatively similar percentage of patients presenting with systolic blood pressure <90 mm Hg (4) and similar admission systolic blood pressure (2) compared with U.S. patients. These management differences may have important prognostic implications, as even short-term inotrope use is associated with increased mortality (32).

Uptake of guideline-directed medical therapy (GDMT) in Asia has exhibited a distinct pattern. In ADHERE-AP, compared with ADHERE, use of angiotensin-converting enzyme inhibitor/angiotensin receptor blocker (ACEI/ARB) was similar, whereas use of mineralocorticoid receptor antagonists was higher in Asia, and use of beta-blockers lower (4). These observations may be related to perceptions of drug tolerability in Asian populations. Importantly, significant regional differences in prescription of GDMT exist even within Asia, with higher rates in Japan (20) and lower use in developing countries (2). Trial datasets such as the ASTRONAUT (Aliskiren Trial on Acute Heart Failure Outcomes) and ASCEND-HF (Acute Study of Clinical Effectiveness of Nesiritide in Decompensated Heart Failure) trials demonstrated similar differences in a comparison of Asia-Pacific to other regions (33,34). Factors associated with underutilization of medications in Asia include rural residency, less-specialized health care providers, and fewer comorbid conditions (35). Analyses have also assessed chronic HF patients of Asian descent who now reside in Western countries. For instance, Chinese individuals with HF living in Canada reported lower use of ACEIs than non-Asians (36). Importantly, data suggest that the use of GDMT in Asia has increased in recent years. For instance, in Japan, ACEI/ARB and beta-blocker use increased from 69% and 28%, respectively, to 72% and 49%, respectively, from 2000 to 2010 (19).

**PHARMACOLOGIC DIFFERENCES.** Few data are available regarding differences in dosing, tolerability, or adherence in Asia compared with other regions. Nonetheless, geographic differences in the efficacy of GDMT may exist (37). Distinct HF phenotypes among Asians support assessment of differences in the pharmacokinetics and pharmacodynamics for GDMT. Previous studies have demonstrated genetic variations between the renin-angiotensin aldosterone system in Chinese and that in Caucasian populations involving polymorphisms in the ACE and angiotensinogen genes (38,39). Several small studies have suggested that ACEIs demonstrate a pharmacologic profile in Asians that is different from that in Caucasians, including differences in volume of distribution and drug clearance as well as effects on renin-angiotensin aldosterone system levels and blood pressure (40,41). Chinese patients may experience more coughing from ACEIs than Caucasians (38). Similarly, ethnic differences in genetics and pharmacologic response for beta-blockers have been identified. For instance, polymorphisms in hepatic metabolizing proteins that are commonly seen in

Asian populations affect beta-blocker concentrations and clearance (42). Given the small sample size of previous studies and the relative paucity of data pertaining to differences in pharmacological response in Asians (43), future research is needed to clarify the clinical relevance of these findings. Perspectives related to differential pharmacological responses in Asians are highlighted in the 2011 Japanese guidelines (44). For 32% (44 of 137) of the drugs approved in Japan between 2001 and 2007, the maximum recommended dose was less than one-half that in the United States (45). For example, the dosage of carvedilol is recommended to be increased up to 100 mg/day, if tolerated, in the United States and Europe, whereas the maximum approved dosage in Japan is 20 mg/day. Without robust pharmacodynamic studies and dose titration studies that document consistent differences among Asian populations, compared with other world regions, whether the differences in dosages are appropriate remains largely unknown.

**DEVICE THERAPY.** ASTRONAUT and ASCEND-HF reported markedly lower use of implantable cardioverter-defibrillators (ICD) in Asia-Pacific than in other regions (33,34). In ASTRONAUT, only 5.7% of Asian-Pacific patients had an ICD compared with 38.2% in North America, despite similar EF and symptom class. CHART-2 (Chronic Heart Failure Analysis and Registry in the Tohoku District) reported that only 6.6% of Asians with reduced left ventricle EF (LVEF) received a primary prevention ICD (46). It has been suggested that limited accessibility and affordability are primary reasons for low implantation (47). Poorly defined sociocultural norms, conservative value systems, and ethnicity- or religion-specific health beliefs may also play a role.

Other considerations include continued controversy regarding the risk of sudden cardiac death (SCD) in Asians. In the United States, the incidence of SCD was reported to be lower among Asian Americans compared with Caucasians (48). A Japanese study found that Asians who were eligible by MADIT-II criteria but did not undergo ICD implantation had significantly lower risk of SCD and even better overall survival than the historical Western MADIT-II population (49). On the other hand, when MADIT-II criteria were applied to a Chinese cohort, those fulfilling criteria were found to be at similar risk of SCD compared with the original Western MADIT-II population (50). Acknowledging that previous studies were limited by retrospective design and referral and selection bias, the ongoing prospective ASIAN-HF study was designed (51).

## OUTCOMES

Regional differences in HF outcomes have been described (33,52-54). In ADHERE-AP, the median hospital length of stay (LOS) for HF was 6 days, and in-hospital mortality was 4.8% compared with 4 days and 4.0%, respectively, in ADHERE. This may be attributed, in part, to the ADHERE-AP cohort having an increased severity of disease due to larger enrollment from tertiary hospitals. In the Japanese ATTEND registry, median LOS was 21 days, and in-hospital mortality was 6.4% (20). Japanese patients often participate in in-patient disease management programs that increase LOS. Differences in HF disease severity, clinical practice patterns, reimbursement, and participation in disease management programs have been hypothesized to at least partially explain these observations. In contrast, trial data from ASTRONAUT demonstrated similar LOS in North America and Asia-Pacific (33), highlighting differences between trial and registry patients.

Limited data are available regarding post-discharge outcomes in Asia. In Korea, 30-day and 180-day all-cause mortality were 1.2% and 9.2%, respectively, whereas HF readmission rates were 6% and 24%, respectively, at those time points (31). These figures are lower than in other regions in Asia. In ASTRONAUT, the 30-day all-cause mortality and HF hospitalization in Asia-Pacific were 2.7% and 12.5%, respectively, with 12-month rates of 26.7% and 25.1%, respectively (33). Mortality rates in Asia-Pacific were higher than in other regions, but hospitalization rates were lower. In ASTRONAUT, Asia-Pacific enrollment location was independently associated with a  $\geq 3$ -fold increase in mortality compared with that in North America. The rate of SCD at 12 months in the Asia-Pacific region (10.3%) was more than double that in any other world region. These observations can be interpreted in the context of a study of 1,719 HF patients in Singapore, where 1-year mortality was 16%, and 4.5% of subjects had an ICD and/or cardiac resynchronization therapy (CRT), which was associated with clinical outcomes (55).

## ASIAN REPRESENTATION IN TRIALS

Prior large-scale outcome trials of current HF with reduced EF (HF<sub>r</sub>EF) GDMT did not routinely enroll Asians. Specifically, none of the landmark ACEI trials (CONSENSUS, SOLVD, SAVE, AIRE, TRACE) or beta-blocker trials (COPERNICUS, CIBIS-II, MERIT-HF) included patients from Asia (5). In contrast, 2 of the 3 MRA trials enrolled patients from Asia. RALES enrolled patients from Japan (56), and EMPHASIS

enrolled in Hong Kong, Korea, India, Singapore, and the United Arab Emirates (57). Similarly, trials assessing ivabradine and LCZ-696 had greater representation from Asian countries. SHIFT enrolled 532 patients (8.2%) in China, India, Malaysia, and South Korea (7), and PARADIGM-HF enrolled 1,509 patients (18%) in China, Philippines, Singapore, South Korea, Taiwan, and Thailand (6). In addition to these trials, which form the basis of GDMT, ASCEND-HF (58) and ASTRONAUT (33) enrolled 1,762 patients (25%) and 439 patients (27%) from Asia-Pacific, respectively. However, earlier large-scale HF trials, including PROTECT (59) and EVEREST (60), did not enroll in Asia.

### ASIA-SPECIFIC STUDIES

Relatively few Asia-specific studies in HF patients have been conducted. MAIN-CHF-II was a randomized trial of bisoprolol versus carvedilol for 32 weeks in 59 Japanese HF<sub>rEF</sub> patients (61). The study had been stopped earlier after off-label use of bisoprolol was approved in Japan. Importantly, the dosing strategy targeted significantly lower doses than the landmark trials with the drugs. Bisoprolol was started at 0.625 mg daily with up-titration to 5 mg versus 1.25 mg with up-titration to 10 mg in CIBIS-II (62). Similarly, carvedilol was started at 2.5 mg/day and up-titrated to 20 mg/day versus 3.125 mg twice daily with up-titration to 25 mg twice daily in COPERNICUS (63).

SUGAR was an observational study of HF<sub>rEF</sub> patients in Korea, which assessed the association between prescription of GDMT and outcomes (64). Patients who were prescribed GDMT tended to have reduced mortality and rehospitalization at 90 days and 12 months compared to similar patients who did not receive GDMT. With a sample size of 1,319 patients, this analysis was likely underpowered, yet consistent benefits were observed for ACEI/ARBs and beta-blockers. Additional observational studies have supported benefits of beta-blockers in elderly Japanese patients (65). Another analysis assessed carvedilol use in Japanese patients with either preserved or reduced EF (66). Similar observational analyses of ACEI/ARBs in HF<sub>rEF</sub> patients have been performed in Asian populations. For instance, the JCARE-CARD investigators demonstrated similar clinical outcomes when comparing ACEI and ARBs in chronic HF<sub>rEF</sub> patients in Japan (67). However, the lack of control groups, the inclusion of reduced and preserved EF patients, and the observational nature of these studies limit interpretation of results.

The ongoing RELAX-AHF-ASIA trial is exclusively enrolling patients in Asia (NCT02007720). This trial is targeting enrollment of 1,520 patients within Asia in parallel to the overall international RELAX-AHF-2 trial with target enrollment of 6800 patients (NCT01870778).

### FUTURE DIRECTIONS

Given the differences in phenotype and medical management as well as potential differences in pharmacological response to HF medications in Asia, we suggest several design considerations for future trials to improve applicability to Asian populations. Importantly, early-phase pharmacokinetic and dose-ranging studies are needed in Asian populations. Subsequently, we propose a developmental strategy whereby trials are conducted in Asian countries simultaneously with efforts in other regions, rather than the historical model where Asian countries were included only after initial efforts in Europe/North America. Patients from Asia may be included as a pre-specified subgroup of trials or in separate region-specific trials.

In the former strategy, trials are designed to include pre-specified targets of numbers and percentages to be enrolled in Asia, in order for meaningful conclusions to be drawn within this pre-specified subgroup. This strategy applies uniform protocols in Asia and elsewhere, with post hoc statistical techniques to adjust for baseline differences and assess for interaction between region/ethnicity and drug response.

In the latter strategy, there may be sufficient interest in international or within-region differences in response to a given compound, or anticipated differences in clinical settings or practice patterns, to warrant a dedicated trial in Asia. Such a trial may run in parallel with trials in the rest of the world, but with the Asian trial uniquely designed to account for Asia-specific patient and practice characteristics. For example, the RELAX-ASIA trial accounted for regional differences in patient management pathways for acute HF, and included region-specific renal function cutoffs. Lower age and BMI cutoffs may also be needed for HF trials in Asia.

Differences in background therapy may influence trial design. As an example, the heart rate-reducing agent ivabradine is approved for use in chronic HF<sub>rEF</sub> patients in sinus rhythm with a heart rate of  $\geq 70$  beats/min on maximally tolerated beta-blocker dosage. Use of the drug is dependent on the interpretation of maximally tolerated beta-blocker dosage, which may vary across world regions with

potentially lower dosing used in Asian countries. Data are needed to clarify tolerability and target dosing of HF drugs in Asian populations. One strategy to address this evidence gap is to develop region-specific databanks of administrative, registry, and trial data. With the rapid growth and evolving health care systems in many of these countries, several of these databases are being developed (68).

HF trial development and conduct must acknowledge that Asia covers a diverse group of nations, each with unique patient, sociocultural, and medical practice backgrounds that may affect trial design. There is a need to collect blood samples to assess potential genetic differences that might alter responsiveness to drugs or devices, as in the ongoing ASIAN-HF registry (51). Moreover, sociocultural norms of aging, health beliefs, and

receptivity to medical intervention differ within and between regions. LOS varies, thus influencing the utility of using 30-day rehospitalization as an outcome measure in trials. Dedicated HF programs and outpatient clinics allow close follow-up in some nations (e.g., Singapore) but not others (e.g., parts of China and India), where it may be challenging to follow patients from rural communities. Processes for ethics approval and requirements for regulatory approval also vary widely and must be taken into consideration for optimal design of trials in Asia.

---

**REPRINT REQUESTS AND CORRESPONDENCE:** Dr. Robert J. Mentz, Duke Clinical Research Institute, 2301 Erwin Road, PO Box 17969, Durham, North Carolina 27715. E-mail: [robert.mentz@duke.edu](mailto:robert.mentz@duke.edu).

## REFERENCES

- Mozaffarian D, Benjamin EJ, Go AS, et al. Heart disease and stroke statistics—2015 update. *Circulation* 2015;131:e29-322.
- Ambrosy AP, Fonarow GC, Butler J, et al. The global health and economic burden of hospitalizations for heart failure: lessons learned from hospitalized heart failure registries. *J Am Coll Cardiol* 2014;63:1123-33.
- Shimokawa H, Miura M, Nochioka K, Sakata Y. Heart failure as a general pandemic in Asia. *Eur J Heart Fail* 2015;17:884-92.
- Atherton JJ, Hayward CS, Wan Ahmad WA, et al. Patient characteristics from a regional multicenter database of acute decompensated heart failure in Asia Pacific. *J Card Fail* 2012;18:82-8.
- Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA guideline for the management of heart failure. *J Am Coll Cardiol* 2013;62:e147-239.
- McMurray JJ, Packer M, Desai AS, et al. Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med* 2014;371:993-1004.
- Swedberg K, Komajda M, Bohm M, et al. Ivabradine and outcomes in chronic heart failure (SHIFT). *Lancet* 2010;376:875-85.
- Glickman SW, McHutchison JG, Peterson ED, et al. Ethical and scientific implications of the globalization of clinical research. *N Engl J Med* 2009;360:816-23.
- China Food and Drug Administration. Regulations for implementation of the Drug Administration Law of the People's Republic of China. Available at: <http://eng.sfda.gov.cn/WS03/CL0767/61640.html>. Accessed September 10, 2015.
- Ng TP, Niti M. Trends and ethnic differences in hospital admissions and mortality for congestive heart failure in the elderly in Singapore, 1991 to 1998. *Heart* 2003;89:865-70.
- Chong AY, Rajaratnam R, Hussein NR, Lip GY. Heart failure in a multiethnic population in Kuala Lumpur, Malaysia. *Eur J Heart Fail* 2003;5:569-74.
- Tseng CH. Clinical features of heart failure hospitalization in younger and elderly patients in Taiwan. *Eur J Clin Invest* 2011;41:597-604.
- National Center for Cardiovascular Diseases, China. Report on cardiovascular diseases in China (2011). In: Hu Shengshou, Kong Lingzhi, editors. Beijing, China: Encyclopedia of China Publishing House, 2014.
- Huffman MD, Prabhakaran D. Heart failure: epidemiology and prevention in India. *Natl Med J India* 2010;23:283-8.
- Seferovic PM, Stoerk S, Filippatos G, et al. Organization of heart failure management in European Society of Cardiology member countries: survey of the Heart Failure Association of the European Society of Cardiology in collaboration with the Heart Failure National Societies/Working Groups. *Eur J Heart Fail* 2013;15:947-59.
- Guo Y, Lip GY, Banerjee A. Heart failure in East Asia. *Curr Cardiol Rev* 2013;9:112-22.
- Singapore Ministry of Health. Heart Failure Costs and Financing. Available at: [https://www.moh.gov.sg/content/moh\\_web/home/costs\\_and\\_financing/HospitalBillSize/heart\\_failure.html](https://www.moh.gov.sg/content/moh_web/home/costs_and_financing/HospitalBillSize/heart_failure.html). Accessed October 1, 2015.
- Yearbook of Statistics Singapore 2014. Department of Statistics Singapore. [www.singstat.gov.sg](http://www.singstat.gov.sg). ISSN 0583-3655. Accessed October 1, 2015.
- Shiba N, Nochioka K, Miura M, Kohno H, Shimokawa H. Trend of Westernization of etiology and clinical characteristics of heart failure patients in Japan—first report from the CHART-2 study. *Circ J* 2011;75:823-33.
- Sato N, Kajimoto K, Keida T, et al. Clinical features and outcome in hospitalized heart failure in Japan (from the ATTEND Registry). *Circ J* 2013;77:944-51.
- Guo Y, Tian Y, Wang H, Si Q, Wang Y, Lip GYH. Prevalence, incidence, and lifetime risk of atrial fibrillation in China. *Chest* 2015;147:109-19.
- Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: the Framingham Study. *Stroke* 1991;22:983-8.
- Colilla S, Crow A, Petkun W, Singer DE, Simon T, Liu X. Estimates of current and future incidence and prevalence of atrial fibrillation in the U.S. adult population. *Am J Cardiol* 2013;112:1142-7.
- Stefansdottir H, Aspelund T, Gudnason V, Arnar DO. Trends in the incidence and prevalence of atrial fibrillation in Iceland and future projections. *Europace* 2011;13:1110-7.
- Tse HF, Wang YJ, Ahmed Ai-Abdullah M, et al. Stroke prevention in atrial fibrillation—an Asian stroke perspective. *Heart Rhythm* 2013;10:1082-8.
- Nguyen CT, Pham NM, Lee AH, Binns CW. Prevalence of and risk factors for type 2 diabetes mellitus in vietnam: a systematic review. *Asia Pac J Public Health* 2015;27:588-600.
- Jan Mohamed HJ, Yap RW, Loy SL, Norris SA, Biesma R, Aagaard-Hansen J. Prevalence and determinants of overweight, obesity, and type 2 diabetes mellitus in adults in Malaysia. *Asia Pac J Public Health* 2015;27:123-35.
- Shi L, Shu XO, Li H, et al. Physical activity, smoking, and alcohol consumption in association with incidence of type 2 diabetes among middle-aged and elderly Chinese men. *PLoS One* 2013;8:e77919.
- IDF Diabetes Atlas 2014. Available at: <http://www.idf.org/diabetesatlas/5e/diabetes>. Accessed August 18, 2015.
- Adams KF Jr., Fonarow GC, Emerman CL, et al. Characteristics and outcomes of patients hospitalized for heart failure in the United States: rationale, design, and preliminary observations from the first 100,000 cases in the Acute Decompensated Heart Failure National Registry (ADHERE). *Am Heart J* 2005;149:209-16.



31. Lee SE, Cho HJ, Lee HY, et al. A multicentre cohort study of acute heart failure syndromes in Korea: rationale, design, and interim observations of the Korean Acute Heart Failure (KorAHF) registry. *Eur J Heart Fail* 2014;16:700-8.
32. Abraham WT, Adams KF, Fonarow GC, et al. In-hospital mortality in patients with acute decompensated heart failure requiring intravenous vasoactive medications: an analysis from the ADHERE. *J Am Coll Cardiol* 2005;46:57-64.
33. Greene SJ, Fonarow GC, Solomon SD, et al. Global variation in clinical profile, management, and post-discharge outcomes among patients hospitalized for worsening chronic heart failure: findings from the ASTRONAUT trial. *Eur J Heart Fail* 2015;17:591-600.
34. Howlett JG, Ezekowitz JA, Podder M, et al. Global variation in quality of care among patients hospitalized with acute heart failure in an international trial: findings from ASCEND-HF. *Circ Cardiovasc Qual Outcomes* 2013;6:534-42.
35. Kim JY, Kim HJ, Jung SY, et al. Utilization of evidence-based treatment in elderly patients with chronic heart failure: using Korean health insurance claims database. *BMC Cardiovasc Disord* 2012;12:60.
36. Choi D, Nemi E, Fernando C, Gupta M, Moe GW. Differences in the clinical characteristics of ethnic minority groups with heart failure managed in specialized heart failure clinics. *JACC Heart Fail* 2014;2:392-9.
37. O'Connor CM, Fiuzat M, Swedberg K, et al. Influence of global region on outcomes in heart failure beta-blocker trials. *J Am Coll Cardiol* 2011;58:915-22.
38. Ding PY, Hu OY, Pool PE, Liao W. Does Chinese ethnicity affect the pharmacokinetics and pharmacodynamics of angiotensin-converting enzyme inhibitors? *J Hum Hypertens* 2000;14:163-70.
39. Thomas GN, Young RP, Tomlinson B, Woo KS, Sanderson JE, Critchley JA. Renin-angiotensin-aldosterone system gene polymorphisms and hypertension in Hong Kong Chinese. *Clin Exp Hypertens* 2000;22:87-97.
40. Anderson PJ, Critchley JA, Tomlinson B. A comparison of the pharmacokinetics and pharmacodynamics of cilazapril between Chinese and Caucasian healthy, normotensive volunteers. *Eur J Clin Pharmacol* 1996;50:57-62.
41. Hu OY, Ding PY, Huang CS, Hwang GM, Chu KM. Pharmacokinetics of fosinoprilat in Chinese and whites after intravenous administration. *J Clin Pharmacol* 1997;37:834-40.
42. Chan SW, Hu M, Ko SS, et al. CYP2C19 genotype has a major influence on labetalol pharmacokinetics in healthy male Chinese subjects. *Eur J Clin Pharmacol* 2013;69:799-806.
43. Ogawa R, Stachnik JM, Echizen H. Clinical pharmacokinetics of drugs in patients with heart failure. *Clin Pharmacokinet* 2014;53:1083-114.
44. Ministry of Health, Labour and Welfare. Guidelines on clinical evaluation of drugs to treat heart failure. Notification No. 0329 (18) of the Evaluation and Licensing Division, PFSB dated March 29, 2011.
45. Shinagawa K. Clinical development and regulatory approval of acute heart failure drugs in Japan. *Cardiovasc Drugs Ther* 2015;29:107-9.
46. Satake H, Fukuda K, Sakata Y, et al. Current status of primary prevention of sudden cardiac death with implantable cardioverter defibrillator in patients with chronic heart failure—a report from the CHART-2 Study. *Circ J* 2015;79:381-90.
47. Pillai HS, Ganapathi S. Heart failure in South Asia. *Curr Cardiol Rev* 2013;9:102-11.
48. Zheng ZJ, Croft JB, Giles WH, Mensah GA. Sudden cardiac death in the United States, 1989 to 1998. *Circulation* 2001;104:2158-63.
49. Tanno K, Miyoshi F, Watanabe N, et al. Are the MADIT II criteria for ICD implantation appropriate for Japanese patients? *Circ J* 2005;69:19-22.
50. Siu CW, Pong V, Ho HH, et al. Are MADIT II criteria for implantable cardioverter defibrillator implantation appropriate for Chinese patients? *J Cardiovasc Electrophysiol* 2010;21:231-5.
51. Lam CS, Anand I, Zhang S, et al. Asian Sudden Cardiac Death in Heart Failure (ASIAN-HF) registry. *Eur J Heart Fail* 2013;15:928-36.
52. Mentz RJ, Kaski JC, Dan GA, et al. Implications of geographical variation on clinical outcomes of cardiovascular trials. *Am Heart J* 2012;164:303-12.
53. Mentz RJ, Cotter G, Cleland JG, et al. International differences in clinical characteristics, management, and outcomes in acute heart failure patients: better short-term outcomes in patients enrolled in Eastern Europe and Russia in the PROTECT trial. *Eur J Heart Fail* 2014;16:614-24.
54. Blair JE, Zannad F, Konstam MA, et al. Continental differences in clinical characteristics, management, and outcomes in patients hospitalized with worsening heart failure results from the EVEREST (Efficacy of Vasopressin Antagonism in Heart Failure: Outcome Study with Tolvaptan) program. *J Am Coll Cardiol* 2008;52:1640-8.
55. Lee JF, Yeo PSD, Koh A, Ho CY, Foo CGD. Implantable Cardioverter-Defibrillator Use to Prevent Sudden Cardiac Death in Eligible Heart Failure Patients is Rare in Singapore. *J Card Fail* 2014;20(85):546.
56. Pitt B, Zannad F, Remme WJ, et al. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. *N Engl J Med* 1999;341:709-17.
57. Zannad F, McMurray JJ, Krum H, et al. Eplerenone in patients with systolic heart failure and mild symptoms. *N Engl J Med* 2011;364:11-21.
58. O'Connor CM, Starling RC, Hernandez AF, et al. Effect of nesiritide in patients with acute decompensated heart failure. *N Engl J Med* 2011;365:32-43.
59. Massie BM, O'Connor CM, Metra M, et al. Roflofylline, an adenosine A1-receptor antagonist, in acute heart failure. *N Engl J Med* 2010;363:1419-28.
60. Konstam MA, Gheorghiade M, Burnett JC Jr., et al. Effects of oral tolvaptan in patients hospitalized for worsening heart failure: the EVEREST Outcome Trial. *JAMA* 2007;297:1319-31.
61. Hori M, Nagai R, Izumi T, Matsuzaki M. Efficacy and safety of bisoprolol fumarate compared with carvedilol in Japanese patients with chronic heart failure: results of the randomized, controlled, double-blind, Multistep Administration of bisoprolol IN Chronic Heart Failure II (MAIN-CHF II) study. *Heart Vessels* 2014;29:238-47.
62. The Cardiac Insufficiency Bisoprolol Study II (CIBIS-II): a randomised trial. *Lancet* 1999;353:9-13.
63. Packer M, Coats AJ, Fowler MB, et al. Effect of carvedilol on survival in severe chronic heart failure. *N Engl J Med* 2001;344:1651-8.
64. Yoo BS, Oh J, Hong BK, et al. Survey of Guideline Adherence for Treatment of Systolic Heart Failure in Real World (SUGAR): a multi-center, retrospective, observational study. *PLoS one* 2014;9:e86596.
65. Yanagihara K, Kinugasa Y, Sugihara S, et al. Discharge use of carvedilol is associated with higher survival in Japanese elderly patients with heart failure regardless of left ventricular ejection fraction. *J Cardiovasc Pharmacol* 2013;62:485-90.
66. Mori Y, Nishikawa Y, Kobayashi F, Hiramatsu K. Clinical status and outcome of Japanese heart failure patients with reduced or preserved ejection fraction treated with carvedilol. *Int Heart J* 2013;54:15-22.
67. Tsuchihashi-Makaya M, Furumoto T, Kinugawa S, et al. Discharge use of angiotensin receptor blockers provides comparable effects with angiotensin-converting enzyme inhibitors on outcomes in patients hospitalized for heart failure. *Hypertension Res* 2010;33:197-202.
68. Kaneko H, Suzuki S, Yajima J, et al. Clinical characteristics and long-term clinical outcomes of Japanese heart failure patients with preserved versus reduced left ventricular ejection fraction: a prospective cohort of Shinken Database 2004-2011. *J Cardiol* 2013;62:102-9.
69. Hamaguchi S, Kinugawa S, Tsuchihashi-Makaya M, et al. Loop diuretic use at discharge is associated with adverse outcomes in hospitalized patients with heart failure: a report from the Japanese cardiac registry of heart failure in cardiology (JCARE-CARD). *Circ J* 2012;76:1920-7.
70. Fonarow GC, Abraham WT, Albert NM, et al. Influence of a performance-improvement initiative on quality of care for patients hospitalized with heart failure: results of the Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients With Heart Failure (OPTIMIZE-HF). *Arch Intern Med* 2007;167:1493-502.
71. O'Connor CM, Abraham WT, Albert NM, et al. Predictors of mortality after discharge in patients hospitalized with heart failure: an analysis from the Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients with Heart Failure (OPTIMIZE-HF). *Am Heart J* 2008;156:662-73.
72. Nieminen MS, Brutsaert D, Dickstein K, et al. EuroHeart Failure Survey II (EHFS II): a survey on hospitalized acute heart failure patients: description of population. *Eur Heart J* 2006;27:2725-36.

**KEY WORDS** Asia, heart failure, trials

**APPENDIX** For supplemental tables, please see the online version of this article.